

Neutrino-cooled Accretion Disks around Spinning Black Holes

Wen-xin Chen* and Andrei M. Beloborodov*

**Physics Department and Columbia Astrophysics Laboratory, Columbia University, 538 West 120th Street, New York, NY 10027*

Abstract. We calculate the structure of accretion disk around a spinning black hole for accretion rates $\dot{M} = 0.01 - 10M_{\odot} s^{-1}$. The model is fully relativistic and treats accurately the disk micro-physics including neutrino emissivity, opacity, electron degeneracy, and nuclear composition. We find that the accretion flow always regulates itself to a mildly degenerate state with the proton-to-nucleon ratio $Y_e \lesssim 0.1$ and becomes very neutron-rich. The disk has a well defined “ignition” radius where neutrino flux raises dramatically, cooling becomes efficient, and Y_e suddenly drops. We also calculate other characteristic radii of the disk, including the ν -opaque and ν -trapping radii, and show their dependence on $\dot{M}$. Accretion disks around fast-rotating black holes produce intense neutrino fluxes which may deposit enough energy above the disk to generate a GRB jet.

Most GRB models assume a short-lived accretion disk as a primary source of energy. Such disks have huge accretion rates $\dot{M} = 0.01 - 10M_{\odot} s^{-1}$ and their cooling mechanism is neutrino emission [1, 2, 3]. Special features of neutrino-cooled disks are:

- The main cooling is due to reactions $p + e^{-} \rightarrow n + \nu$ and $n + e^{+} \rightarrow p + \bar{\nu}$.
- Electron degeneracy affects the disk structure.
- The disk can be opaque to neutrinos which affects the cooling rate.
- Nuclear composition changes with radius, consuming energy.
- Neutrino cooling and electron degeneracy lead to very high neutron richness.

The disk model may be formulated following the classical vertically-integrated approach of Shakura & Sunyaev and approximating the azimuthal motion in the disk, u^{ϕ} , as Keplerian rotation. A small radial velocity is superimposed on this rotation. It is approximately given by $u^r = \alpha B(H/r)^2 r u^{\phi}$ where H is half-thickness of the disk (found from hydrostatic balance), B is a numerical factor determined by Kerr metric, and $\alpha = 0.1 - 0.01$.

The main equations of the accretion disk are as follows.

$$\text{Energy conservation :} \quad F^{+} - F^{-} = cu^r \left[\frac{d(UH)}{dr} - \frac{U+P}{\rho} \frac{d(\rho H)}{dr} \right]. \quad (1)$$

$$\text{Lepton conservation :} \quad \frac{1}{H} (\dot{N}_{\bar{\nu}} - \dot{N}_{\nu}) = cu^r \left[\frac{\rho}{m_p} \frac{dY_e}{dr} + \frac{d}{dr} (n_{\nu} - n_{\bar{\nu}}) \right]. \quad (2)$$

$$\text{Charge conservation :} \quad n_{e^{-}} - n_{e^{+}} = Y_e \frac{\rho}{m_p}. \quad (3)$$

$$\text{Chemical balance :} \quad \mu_e - \mu_{\nu} = kT \ln \left(\frac{1 - Y_e}{Y_e} \right) + (m_n - m_p) c^2. \quad (4)$$

Here F^+ is the viscously dissipated energy per unit area, F^- is the neutrino energy flux, $\dot{N}_\nu$ and $\dot{N}_{\bar{\nu}}$ are the number fluxes of neutrinos and antineutrinos, ρ is the mass density, T is temperature, Y_e is the proton-to-baryon ratio, μ_e and μ_ν are the chemical potentials of electrons and neutrinos, n_{e^-} and n_{e^+} are number densities of electrons and positrons. The energy density U and pressure P include contributions of baryons, electrons, positrons, radiation, and neutrinos. They are calculated by integrating the corresponding distribution functions (Fermi-Dirac distributions for $e^\pm$ and $\nu, \bar{\nu}$ if the disk is opaque to neutrinos).

Note that we include the advection terms in the laws of energy and lepton conservation (right-hand side of eqs. 1 and 2). They describe the most important effects of advection. Other effects of heat storage in the disk — in particular, the dynamical effects of the radial pressure gradient on u^r and u^ϕ — are relatively small and neglected here, i.e. the standard α -disk model is used to calculate u^ϕ and u^r .

The chemical balance and μ_ν are used only in the ν -opaque region; then the cooling rate is calculated as $F^- = cU_\nu/\tau_\nu + cU_{\bar{\nu}}/\tau_{\bar{\nu}}$. In the ν -transparent region, F^- equals $2H(\dot{U}_{e^-p} + \dot{U}_{e^+n})$ where $\dot{U}_{e^-p}$ and $\dot{U}_{e^+n}$ are the rates of energy release by reactions $e^- + p \rightarrow n + \nu$ and $e^+ + n \rightarrow p + \bar{\nu}$. The abundance of α -particles is determined by the nuclear statistical equilibrium.

We have solved the set of disk equations numerically starting from an outer region where no neutrino cooling takes place and assuming that the infalling matter is initially made of α particles (heavier elements are decomposed into α -particles as the matter approaches the black hole). A schematic picture of the accretion disk and its characteristic radii are shown in Figure 1.




FIGURE 1. Schematic picture of the disk with characteristic radii indicated.

The outer region $r > 50r_g$ ($r_g = 2GM/c^2$ is the gravitational radius of the black hole) is dominated by α particles. Neutrino cooling is negligible in this region and the viscously dissipated heat is stored in the disk and advected inward. The destruction of α particles at $r_\alpha \sim 50r_g$ consumes ~ 7 MeV per baryon, which makes the disk somewhat thinner. At the "ignition" radius r_{ign} (slightly smaller than r_α), kT reaches $\sim$ MeV and neutrino cooling becomes significant, further reducing disk scale-height H/r . With decreasing radius the disk becomes opaque to neutrinos (first to ν and then to $\bar{\nu}$). In the innermost region of disks with $\dot{M} > 2M_\odot \text{ s}^{-1}$ the neutrinos are trapped and advected toward the black hole. Figure 2 shows the found characteristic radii as functions of $\dot{M}$ for disks with $\alpha = 0.1$ around black holes with spin parameters $a = 0$ and 0.95 .

We find two general properties of the disk:

1. — It regulates itself to a mildly degenerate state with $\mu_e = 1 \sim 3kT$. The reason of this regulation is the negative feedback of degeneracy and neutronization on the cooling rate: higher




FIGURE 2. Characteristic radii in the accretion disk as a function of $\dot{M}$ for a black hole of mass $M = 3M_\odot$ and spin parameter $a = 0.95$ (left) and 0 (right). Viscosity parameter $\alpha = 0.1$ is assumed. Neutrino cooling does not ignite when $\dot{M}$ is below $0.02M_\odot s^{-1}$ for Kerr and $0.1M_\odot s^{-1}$ for Schwarzschild black holes. Neutrino trapping takes place for Kerr black holes when $\dot{M} > 2M_\odot s^{-1}$. The disk extends down to the marginally stable orbit, which is $3r_g$ for $a = 0$ and $\approx r_g$ for $a = 0.95$.

degeneracy $\mu_e/kT \rightarrow$ fewer positrons ($n_+/n_- \sim e^{-\mu_e/kT}$) $\rightarrow$ lower equilibrium rate of neutrino emission $\rightarrow$ lower cooling rate $\rightarrow$ higher temperature $\rightarrow$ lower degeneracy.

2. — In a broad range of relevant accretion rates the disk is found to reach $Y_e \sim 0.1$. It corresponds to a very high neutron-to-proton ratio ~ 10 .

Figure 3 shows examples of the disk structure around a Kerr black hole with $a = 0.95$ with accretion rate $0.2M_\odot s^{-1}$, for three different values of viscosity parameter α . The transitions at the characteristic radii manifest themselves in the non-monotonic behavior of the degeneracy parameter $\eta = \mu_e/kT$, F^-/F^+ , Y_e , and H/r . In particular the sharp maximum of F^-/F^+ appears at the ignition radius. The disk is quite thin at $r < r_{ign}$ and cooled locally, $F^- \approx F^+$.

The high neutron richness has important implications for the global picture of GRB explosion. When the neutron-rich material is ejected in a relativistic jet and develops a large Lorentz factor, the neutrons decay at distance $\sim 10^{16}$ cm and affect the observed explosion.

Disks around rapidly rotating black holes produce a high neutrino flux. It deposits energy above the disk (via neutrino annihilation $\nu + \bar{\nu} \rightarrow e^+ + e^-$) and can drive an outflow with power $L > 10^{51}$ erg/s. A detailed study of $\nu\bar{\nu}$ annihilation around a Kerr black hole is in preparation.

ACKNOWLEDGMENTS

This work was supported by NASA grant NAG5-13382.

REFERENCES

1. R. Popham, S.E. Woosley, & C. Fryer, *Astrophys. J.*, **518**, 356 (1999).
2. A.M. Beloborodov, *Astrophys. J.*, **588**, 931 (2003).
3. K. Kohri, R. Narayan, & T. Piran, *Astrophys. J.*, **629**, 341 (2005).




FIGURE 3. Numerical solution for accretion disk with $\dot{M} = 0.2M_\odot s^{-1}$ around a black hole with mass $M = 3M_\odot$ and spin parameter $a = 0.95$. Three models are shown with viscosity parameter $\alpha = 0.1, 0.03$ and 0.01 . The degeneracy parameter η is defined as μ_e/kT , X_{free} is the mass fraction of free nucleons.